

# HOMOLOGICAL MIRROR SYMMETRY FOR RABINOWITZ FUKAYA CATEGORIES OF MILNOR FIBERS OF BRIESKORN–PHAM SINGULARITIES

YANKI LEKILI AND KAZUSHI UEDA

ABSTRACT. We discuss homological mirror symmetry for Rabinowitz Fukaya categories of Milnor fibers of invertible polynomials, and prove it for Brieskorn–Pham polynomials which are not of Calabi–Yau type. This allows a calculation of the Rabinowitz Floer homology of the Milnor fiber as the Hochschild homology of the dg category of equivariant matrix factorizations.

## 1. INTRODUCTION

1.1. For a pair  $(k, m)$  of integers satisfying  $k \geq 2$  and  $m - k \geq 2$ , let

$$(1.1) \quad \check{U} := \{(x, y, z, w) \in \mathbb{C}^4 \mid x^k + y^{m-k} + z^2 + w^2 = 1\}$$

be the Milnor fiber of the double suspension of the Brieskorn–Pham polynomial  $x^k + y^{m-k}$ . The cases  $(k, m) = (2, r + 3)$ ,  $(3, 6)$ ,  $(3, 7)$ , and  $(3, 8)$  give simple singularities of types  $A_r$ ,  $D_4$ ,  $E_6$ , and  $E_8$  respectively, for whom homological mirror symmetry conjectured in [LU22] is proved in [LU21].

1.2. Fix an algebraically closed field  $\mathbf{k}$  of characteristic zero as coefficients for Fukaya categories. The *stable Fukaya category* of the Liouville manifold  $\check{U}$  is defined as the quotient

$$(1.2) \quad \mathcal{S}(\check{U}) := \mathcal{W}(\check{U}) / \mathcal{F}(\check{U})$$

of the wrapped Fukaya category  $\mathcal{W}(\check{U})$  by its full subcategory  $\mathcal{F}(\check{U})$  consisting of compact Lagrangian submanifolds.

1.3. Set

$$(1.3) \quad R := \mathbf{k}[x, y] / (x^k + y^{m-k}),$$

and let  $G$  be the diagonal subgroup of  $\mathrm{SL}_2$  isomorphic to  $\boldsymbol{\mu}_m = \mathrm{Spec} \mathbf{k}[\xi] / (\xi^m - 1)$ . The stable category  $\underline{\mathrm{CM}}_G(R)$  of the category  $\mathrm{CM}_G(R)$  of  $G$ -equivariant maximal Cohen–Macaulay  $R$ -modules is the homotopy category of the dg category  $\mathrm{mf}([\mathbb{A}^2/G], x^k + y^{m-k})$  of  $G$ -equivariant matrix factorizations of  $x^k + y^{m-k}$  on  $\mathbb{A}^2 = \mathrm{Spec} \mathbf{k}[x, y]$  [Eis80], which is quasi-equivalent to the *stable derived category*  $\mathrm{scoh} X := \mathrm{coh} X / \mathrm{perf} X$  defined as the dg quotient of the bounded derived category of coherent sheaves on the quotient stack  $X := [\mathrm{Spec} R/G]$  by the full subcategory consisting of perfect complexes [Buc87, Orl04].

1.4. As shown in [JKS16], the category  $\mathrm{CM}_G(R)$  gives an additive categorification of the cluster algebra  $\mathbf{k}[\mathrm{Gr}_{k,m}]$ . Moreover, the endomorphism ring of a cluster-tilting module is described by a dimer model on a disk [BKM16], which is originally introduced in [Pos] to describe parametrizations of cells in totally nonnegative Grassmannians, and used in [Sco06] to give a structure of a cluster algebra on the homogeneous coordinate ring of  $\mathrm{Gr}(k, m)$ .

1.5. In this paper, we give two proofs of an equivalence

$$(1.4) \quad \mathrm{mf}([\mathbb{A}^2/G], x^k + y^{m-k}) \simeq \mathcal{S}(\check{U})$$

and its generalizations. From (1.4) and on, dg categories of matrix factorizations and Fukaya categories are completed with respect to cones and direct summands, so that they are idempotent-complete stable  $\infty$ -categories over  $\mathbf{k}$ .

1.5.1. One proof is based on homological mirror symmetry for (exact symplectic Lefschetz fibrations associated with) invertible polynomials, which is known for Brieskorn–Pham singularities [FU11].

1.5.2. The other proof is based on Conjecture 3.2, which is homological mirror symmetry for Milnor fibers of invertible polynomials [LU22]. We prove Conjecture 3.2 for Brieskorn–Pham singularities in Theorem 4.1.

1.6. Let  $\mathrm{RFH}_*(\check{U})$  and  $\mathcal{R}(\check{U})$  be the Rabinowitz Floer homology and the Rabinowitz Fukaya category of  $\check{U}$  respectively. Koszul duality holds for  $\check{U}$  by [LU22, Theorem 6.11], so that one has

$$(1.5) \quad \mathcal{S}(\check{U}) \simeq \mathcal{R}(\check{U})$$

by [GGV, Corollary 1.4] (cf. also [KS]), and

$$(1.6) \quad \mathrm{RFH}_*(\check{U}) \simeq \mathrm{HH}_*(\mathcal{R}(\check{U}))$$

by [GGV, Corollary 1.7]. Hence one can compute the Rabinowitz Floer homology as the Hochschild homology of the dg category of equivariant matrix factorizations.

1.7. This paper is organized as follows:

1.7.1. In Section 2, we prove Theorem 2.2, which shows that a generalization of (1.4) to double suspensions of invertible polynomials follows from Conjecture 2.1. Conjecture 2.1 is homological mirror symmetry for invertible polynomials. Theorem 2.2 implies (1.4) since Conjecture 2.1 is known for Brieskorn–Pham polynomials.

1.7.2. In Section 3, we show that Conjecture 3.2 (which is homological mirror symmetry for Milnor fibers of invertible polynomials) implies (3.19) (which is a ‘stable’ version of homological mirror symmetry for Milnor fibers of invertible polynomials). Then we prove Theorem 3.3 (which shows that a generalization of (1.4) to double suspensions of invertible polynomials follows from Conjecture 3.2) using (3.19) and the Knörrer periodicity.

1.7.3. *Remark.* Theorem 2.2 and Theorem 3.3 have slightly different (but closely related) hypotheses (homological mirror symmetry for  $\mathbf{w}$  for the former and that for the Milnor fiber of  $\mathbf{W} = \mathbf{w} + z^2 + w^2$  for the latter) and the same conclusion.

1.7.4. In Section 4, we prove Conjecture 3.2 for Brieskorn–Pham polynomials. The proof is based on

- a deformation-theoretic argument going back to Seidel and Sheridan, and
- the Koszul duality between the Fukaya category and the wrapped Fukaya category.

1.7.5. The ‘stable’ homological mirror symmetry (3.19) gives an algorithm to compute the Rabinowitz Floer homology explicitly. We give sample calculations in Section 5.

## 2. DOUBLE SUSPENSIONS OF INVERTIBLE POLYNOMIALS

A weighted homogeneous polynomial  $\mathbf{w} \in \mathbb{C}[x_1, \dots, x_n]$  with an isolated critical point at the origin is *invertible* if there is an integer matrix  $A = (a_{ij})_{i,j=1}^n$  with non-zero determinant such that

$$(2.1) \quad \mathbf{w} = \sum_{i=1}^n \prod_{j=1}^n x_j^{a_{ij}}.$$

The *transpose* of  $\mathbf{w}$  is defined in [BH93] as

$$(2.2) \quad \check{\mathbf{w}} = \sum_{i=1}^n \prod_{j=1}^n x_j^{a_{ji}},$$

whose exponent matrix  $\check{A}$  is the transpose matrix of  $A$ . The group

$$(2.3) \quad \Gamma := \{(t_1, \dots, t_n) \in (\mathbb{G}_m)^n \mid t_1^{a_{11}} \cdots t_n^{a_{1n}} = \cdots = t_1^{a_{n1}} \cdots t_n^{a_{nn}}\}$$

acts naturally on  $\mathbb{A}^n$ . The group  $\widehat{\Gamma} := \mathrm{Hom}(\Gamma, \mathbb{G}_m)$  of characters of  $\Gamma$  is generated by  $\chi_i: (t_j)_{j=1}^n \mapsto t_i$  for  $i = 1, \dots, n$  with relations  $\chi := \sum_{i=1}^n a_{1i}\chi_i = \cdots = \sum_{i=1}^n a_{ni}\chi_i$ . Here, the group structure on  $\widehat{\Gamma}$  is written additively.

Let  $\text{mf}([\mathbb{A}^n/\Gamma], \mathbf{w})$  be the idempotent completion of the dg category of  $\Gamma_{\mathbf{w}}$ -equivariant matrix factorizations of  $\mathbf{w}$ , and  $\mathcal{F}(\check{\mathbf{w}})$  be the Fukaya–Seidel category of (a Morsification of)  $\check{\mathbf{w}}$ .

**Conjecture 2.1.** *For any invertible polynomial  $\mathbf{w}$ , one has an equivalence*

$$(2.4) \quad \text{mf}([\mathbb{A}^n/\Gamma], \mathbf{w}) \simeq \mathcal{F}(\check{\mathbf{w}})$$

of  $\infty$ -categories.

Conjecture 2.1 is stated for Brieskorn–Pham singularities in three variables in [Ued06], for polynomials in three variables associated with a regular system of weights of dual type in the sense of Saito in [Tak10], and for invertible polynomials in three variables in [ET11]. It is proved for  $n = 2$  in [HS20], and for Sebastiani–Thom sums of polynomials of type A and D in [FU11, FU13]. The conjecture that  $\text{mf}([\mathbb{A}^n/\Gamma], \mathbf{w})$  has a full exceptional collection, which is implied by Conjecture 2.1, is stated in [HO23, Conjecture 1.4], and proved in [FKK23].

Let  $(\check{d}_1, \dots, \check{d}_n, \check{h})$  be the sequence of positive integers such that  $\gcd(\check{d}_1, \dots, \check{d}_n, \check{h}) = 1$  and

$$(2.5) \quad \check{\mathbf{w}}(t^{\check{d}_1}x_1, \dots, t^{\check{d}_n}x_n) = t^{\check{h}}\check{\mathbf{w}}(x_1, \dots, x_n),$$

which is unique since  $\check{\mathbf{w}}$  is an invertible polynomial. We say that  $\check{\mathbf{w}}$  is of Calabi–Yau type if

$$(2.6) \quad \check{h} = \check{d}_1 + \dots + \check{d}_n.$$

Let

$$(2.7) \quad \check{U} := \{(x_1, \dots, x_n, z, w) \in \mathbb{C}^{n+2} \mid \check{\mathbf{w}} + z^2 + w^2 = 1\}$$

be the Milnor fiber of the double suspension of  $\check{\mathbf{w}}$  and set

$$(2.8) \quad G := \{(t_1, \dots, t_n) \in \Gamma \mid t_1 \cdots t_n = 1\}.$$

We say that  $\mathcal{F}(\check{U})$  and  $\mathcal{W}(\check{U})$  are *Koszul dual* to each other if there exist collections  $(S_i)_{i=1}^{\mu}$  and  $(L_i)_{i=1}^{\mu}$  of objects generating  $\mathcal{F}(\check{U})$  and  $\mathcal{W}(\check{U})$  such that

$$(2.9) \quad \dim_{\mathbf{k}} \text{hom}^*(L_i, S_j) = \delta_{ij}, \quad 1 \leq i, j \leq \mu,$$

where  $\delta_{ij}$  is the Kronecker delta, so that the augmented endomorphism  $A_{\infty}$ -algebras of  $\bigoplus_{i=1}^{\mu} S_i$  and  $\bigoplus_{i=1}^{\mu} L_i$  are Koszul dual to each other. By [LU22, Theorem 6.11], this assumption is satisfied if  $\check{\mathbf{w}}$  is a Brieskorn–Pham polynomial not of Calabi–Yau type.

**Theorem 2.2.** *Conjecture 2.1 and Koszul duality between  $\mathcal{F}(\check{U})$  and  $\mathcal{W}(\check{U})$  implies an equivalence*

$$(2.10) \quad \text{mf}([\mathbb{A}^n/G], \mathbf{w}) \simeq \mathcal{S}(\check{U})$$

of  $\infty$ -categories.

Theorem 2.2 can be regarded as a ‘stable’ homological mirror symmetry for  $\check{U}$ . Theorem 2.2 implies (1.4) since Conjecture 2.1 and Koszul duality are known in this case.

The *n-cluster category* of a pretriangulated  $A_{\infty}$ -category  $\mathcal{A}$  with a Serre functor  $\mathbb{S}$  is defined as the orbit category with respect to the shift  $\mathbb{S}[-n]$  of the Serre functor (see e.g. [Iya18] and references therein). (2.10) is obtained as the composite of equivalences

$$(2.11) \quad \text{mf}([\mathbb{A}^n/G], \mathbf{w}) \simeq C_n(\text{mf}([\mathbb{A}^n/\Gamma], \mathbf{w}))$$

and

$$(2.12) \quad C_n(\mathcal{F}(\check{\mathbf{w}})) \simeq \mathcal{S}(\check{U}).$$

The relation between stable Fukaya categories and cluster categories was first pointed out in [LU22] and studied further in [BJK].

*Proof of (2.11).* Note that one has an isomorphism

$$(2.13) \quad (\chi) \simeq [2]$$

of endofunctors of  $\text{mf}([\mathbb{A}^n/\Gamma], \mathbf{w})$ . Graded Auslander–Reiten duality [AR87] shows that

$$(2.14) \quad \mathbb{S} := (\chi - \chi_1 - \dots - \chi_n)[n - 2]$$

is a Serre functor of  $\text{mf}([\mathbb{A}^n/\Gamma], \mathbf{w})$  (see [IT13, Theorem 2.5]). Hence one has

$$(2.15) \quad \mathbb{S}[-n] \simeq (-\chi_1 - \cdots - \chi_n),$$

so that the orbit category of  $\text{mf}([\mathbb{A}^n/\Gamma], \mathbf{w})$  with respect to  $\mathbb{S}[-n]$  is equivalent to the category of matrix factorizations of  $\mathbf{w}$  graded by  $\widehat{\Gamma}/(-\chi_1 - \cdots - \chi_n) \simeq \widehat{G}$ , which is nothing but the category  $\text{mf}([\mathbb{A}^n/G], \mathbf{w})$  of  $G$ -equivariant matrix factorizations of  $\mathbf{w}$ . This concludes the proof of (2.11).  $\square$

*Proof of (2.12).* Let  $\mathcal{G}$  be the  $(n+1)$ -Calabi–Yau completion of  $\mathcal{A} := \mathcal{F}(\mathbf{w})$  in the sense of [Kel11]. Let further

$$(2.16) \quad \mathcal{B} := \mathcal{A} \oplus \mathcal{A}^\vee[-n-1]$$

be the trivial extension algebra of degree  $n+1$  of  $\mathcal{A}$ , where the  $\mathcal{A}$ -bimodule

$$(2.17) \quad \mathcal{A}^\vee := \text{hom}_{\mathbf{k}}(\mathcal{A}, \mathbf{k})$$

is the graph of the Serre functor. Then  $\mathcal{G}$  is Koszul dual to  $\mathcal{B}$  by [HLW23, Theorem 6]. It follows from [Kel05, Theorem 2] that

$$(2.18) \quad \text{pseu } \mathcal{B} / \text{perf } \mathcal{B} \simeq C_n(\mathcal{A}),$$

where  $\text{pseu } \mathcal{B}$  is the category of *pseudo-perfect*  $\mathcal{B}$ -modules (i.e., dg modules over  $\mathcal{B}$  which are perfect as  $\mathbf{k}$ -modules).

One has

$$(2.19) \quad \mathcal{F}(\check{U}) \simeq \text{perf } \mathcal{B}$$

by [Sei10, Corollary 6.5]. Since  $\mathcal{F}(\check{U})$  and  $\mathcal{W}(\check{U})$  are Koszul dual to each other, one has

$$(2.20) \quad \mathcal{W}(\check{U}) \simeq \text{perf } \mathcal{G}.$$

The Koszul duality between  $\mathcal{G}$  and  $\mathcal{B}$  implies

$$(2.21) \quad \text{pseu } \mathcal{B} \simeq \text{perf } \mathcal{G},$$

and (2.12) is proved.  $\square$

### 3. MORE GENERAL INVERTIBLE POLYNOMIALS

Let

$$(3.1) \quad \check{U} := \{(x_1, \dots, x_{n+2}) \in \mathbb{C}^{n+2} \mid \check{\mathbf{W}} = 1\}$$

be the Milnor fiber of an invertible polynomial

$$(3.2) \quad \check{\mathbf{W}} = \sum_{i=1}^{n+2} \prod_{j=1}^{n+2} x_j^{a_{ji}}$$

in  $n+2$  variables. The group

$$(3.3) \quad K := \{(t_0, \dots, t_{n+2}) \in (\mathbb{G}_m)^{n+3} \mid t_1^{a_{11}} \cdots t_n^{a_{1,n+2}} = \cdots = t_1^{a_{n+2,1}} \cdots t_n^{a_{n+2,n+2}} = t_0 \cdots t_{n+2}\}$$

acts diagonally on  $\mathbb{A}^{n+3}$  making  $\mathbf{W} - x_0 \cdots x_{n+2}: \mathbb{A}^{n+3} \rightarrow \mathbb{A}^1$  equivariant, where

$$(3.4) \quad \mathbf{W} = \sum_{i=1}^{n+2} \prod_{j=1}^{n+2} x_j^{a_{ij}}$$

is the transpose of  $\check{\mathbf{W}}$ . Set  $\check{U} := \text{Spec } \mathbf{k}[x_0, \dots, x_{n+2}]/(\mathbf{W} - x_0 \cdots x_{n+2})$  and  $U := [\check{U}/K]$ , so that

$$(3.5) \quad \text{mf}([\mathbb{A}^{n+3}/K], \mathbf{W} - x_0 \cdots x_{n+2}) \simeq \text{scoh } U.$$

**Lemma 3.1.** *The singular locus of  $\check{U}$  is the  $x_0$ -axis.*

*Proof.* The singular locus of  $\tilde{U}$  is defined inside the ambient space  $\mathbb{A}^{n+3} = \text{Spec } \mathbf{k}[x_0, \dots, x_{n+2}]$  by

$$(3.6) \quad \mathbf{W} - x_0 \cdots x_{n+2} = x_1 \cdots x_{n+2} = \frac{\partial \mathbf{W}}{\partial x_1} - x_0 x_2 \cdots x_{n+2} = \cdots = \frac{\partial \mathbf{W}}{\partial x_{n+2}} - x_0 \cdots x_{n+1} = 0.$$

It follows from  $x_1 \cdots x_{n+2} = 0$  that  $x_i = 0$  for some  $i \in \{1, \dots, n+2\}$ , and one may assume  $i = 1$  without loss of generality. Then one has

$$(3.7) \quad \frac{\partial \mathbf{W}}{\partial x_2} = \cdots = \frac{\partial \mathbf{W}}{\partial x_{n+2}} = 0,$$

and one can show using the classification of invertible polynomials [KS92] that (3.7) implies  $x_j = 0$  for some  $j \in \{2, \dots, n+2\}$ . Then one has

$$(3.8) \quad \frac{\partial \mathbf{W}}{\partial x_1} = x_0 x_2 \cdots x_{n+2} = 0,$$

which together with (3.7) implies  $x_1 = \cdots = x_{n+2} = 0$  since  $\mathbf{W}$  is an invertible polynomial.  $\square$

If  $\mathbf{W}$  is not of Calabi–Yau type, then any point of  $\tilde{U}$  satisfying  $x_0 \neq 0$  can be brought to  $x_0 = 1$  by the action of  $K$ , so that the complement  $D := U \setminus E$  of the closed substack  $E$  of  $U$  defined by  $x_0 = 0$  can be identified with  $[\tilde{D}/H]$  where

$$(3.9) \quad \tilde{D} := \text{Spec } \mathbf{k}[x_1, \dots, x_{n+2}]/(\mathbf{W} - x_1 \cdots x_{n+2})$$

and

$$(3.10) \quad H := \{(t_1, \dots, t_{n+2}) \in (\mathbb{G}_m)^{n+2} \mid t_1^{a_{11}} \cdots t_n^{a_{1,n+2}} = \cdots = t_1^{a_{n+2,1}} \cdots t_n^{a_{n+2,n+2}} = t_1 \cdots t_{n+2}\}$$

$$(3.11) \quad \cong K \cap \{t_0 = 1\}.$$

Since the intersection of the singular locus of  $U$  with  $E$  is the origin, the full subcategory  $\text{scoh}_E U$  of  $\text{scoh } U := \text{coh } U / \text{perf } U$  consisting of objects supported on  $E$  is equivalent to the full subcategory  $\text{scoh}_0 U$  consisting of objects supported at the origin;

$$(3.12) \quad \text{scoh}_0 U \simeq \text{scoh}_E U.$$

**Conjecture 3.2.** *For any invertible polynomial  $\mathbf{W}$  not of Calabi–Yau type, one has equivalences*

$$(3.13) \quad \text{scoh } U \simeq \mathcal{W}(\check{U})$$

and

$$(3.14) \quad \text{scoh}_0 U \simeq \mathcal{F}(\check{U})$$

of  $\infty$ -categories.

(3.13) is [LU22, Conjecture 1.4], from which (3.14) should follow as the restriction to the full subcategories consisting of objects  $X$  such that  $\text{hom}(X, Y)$  is perfect as a  $\mathbf{k}$ -module for any  $Y$ . [Hab22, Theorem 1.1] gives (3.14) for  $n = 0$ , and a  $\mathbb{Z}/2\mathbb{Z}$ -graded variant of (3.13) is discussed in [Gam24].

**Theorem 3.3.** *If Conjecture 3.2 holds for the double suspension of an invertible polynomial  $\mathbf{w}$ , then one has an equivalence*

$$(3.15) \quad \text{mf}([\mathbb{A}^n/G], \mathbf{w}) \simeq \mathcal{S}(\check{U})$$

of  $\infty$ -categories.

*Proof.* The equivalence

$$(3.16) \quad \text{coh } D \simeq \text{coh } U / \text{coh}_E U$$

induces an equivalence

$$(3.17) \quad \text{scoh } D \simeq \text{scoh } U / \text{scoh}_E U,$$

which together with (3.12) gives an equivalence

$$(3.18) \quad \text{scoh } D \simeq \text{scoh } U / \text{scoh}_0 U.$$

It follows that if  $\mathbf{W}$  is an invertible polynomial not of Calabi–Yau type such that Conjecture 3.2 holds, then one has an equivalence

$$(3.19) \quad \text{scoh } D \simeq \mathcal{S}(\check{U})$$

of  $\infty$ -categories.

If  $\mathbf{W}$  is not of Calabi–Yau type, then the singular locus of  $\tilde{D}$  is the origin. Indeed, the singular locus of  $\tilde{D}$  is defined inside the ambient space  $\mathbb{A}^{n+2} = \text{Spec } \mathbf{k}[x_1, \dots, x_{n+2}]$  by

$$(3.20) \quad \mathbf{W} - x_1 \cdots x_{n+2} = \frac{\partial \mathbf{W}}{\partial x_1} - x_2 \cdots x_{n+2} = \cdots = \frac{\partial \mathbf{W}}{\partial x_{n+2}} - x_1 \cdots x_{n+1} = 0.$$

By multiplying  $d_i x_i$  and summing over  $i$ , one obtains

$$(3.21) \quad h\mathbf{W} - \sum_{i=1}^{n+2} d_i x_1 \cdots x_{n+2} = 0,$$

which together with  $\mathbf{W} - x_1 \cdots x_{n+2} = 0$  implies  $\mathbf{W} = x_1 \cdots x_{n+2} = 0$ . Now we can argue as in the proof of Lemma 3.1 that  $x_1 = \cdots = x_{n+2} = 0$ .

The double suspension  $\mathbf{w}(x_1, \dots, x_n) + x_{n+1}^2 + x_{n+2}^2$  is not of Calabi–Yau type since  $2d_{n+1} = 2d_{n+2} = h$  and hence  $\sum_{i=1}^{n+2} d_i > h$ .

Set  $D' := [\tilde{D}'/H]$  where  $\tilde{D}' := \text{Spec } \mathbf{k}[x_1, \dots, x_{n+2}]/(\mathbf{W})$ . The singular loci of both  $\tilde{D}$  and  $\tilde{D}'$  are the origin, and the formal completions of  $\tilde{D}$  and  $\tilde{D}'$  at the origin are isomorphic since

$$(3.22) \quad \mathbf{w}(x_1, \dots, x_n) + x_{n+1}^2 + x_{n+2}^2 - x_1 \cdots x_{n+2} \\ = \mathbf{w}(x_1, \dots, x_n) + \left( \sqrt{1 - \frac{1}{4}(x_1 \cdots x_n)^2 x_{n+1}} \right)^2 + \left( x_{n+2} - \frac{1}{2} x_1 \cdots x_{n+1} \right)^2$$

in  $\mathbf{k}[[x_1, \dots, x_{n+2}]]$ . It follows that

$$(3.23) \quad \text{scoh } D \simeq \text{scoh } D'$$

by [Orl11, Theorem 2.10]. The isomorphism

$$(3.24) \quad H \cong G \times \boldsymbol{\mu}_2 \times \boldsymbol{\mu}_2$$

and the Knörrer periodicity [Knö87, Proposition 2.1] shows

$$(3.25) \quad \text{scoh } D' \simeq \text{mf}([\mathbb{A}^n/G], \mathbf{w}),$$

and Theorem 2.2 is proved.  $\square$

#### 4. HOMOLOGICAL MIRROR SYMMETRY FOR MILNOR FIBERS OF BRIESKORN–PHAM SINGULARITIES

We use the same notations as in Section 3. We prove the following theorem in this section:

**Theorem 4.1.** *Conjecture 3.2 holds if  $\mathbf{W}$  is a Brieskorn–Pham polynomial.*

Let  $M$  be the free abelian group generated by  $\{\mathbf{e}_i\}_{i=1}^{n+2}$ , and  $\tilde{M}$  be the subgroup generated by  $\{\mathbf{f}_j := \sum_{i=1}^{n+2} a_{ji} \mathbf{e}_i\}_{j=1}^{n+2}$ . We write the inclusion  $\tilde{M} \hookrightarrow M$  as  $\varphi$ .

Let

$$(4.1) \quad \mathcal{P} := \{(y_1, \dots, y_{n+2}) \in M_{\mathbb{C}^\times} := M \otimes \mathbb{C}^\times \mid y_1 + \cdots + y_{n+2} + 1 = 0\}$$

be an  $n$ -dimensional pair of pants, and set  $\tilde{\mathcal{P}} := \varphi_{\mathbb{C}^\times}^{-1}(\mathcal{P})$  where

$$(4.2) \quad \varphi_{\mathbb{C}^\times} := \varphi \otimes \mathbb{C}^\times : \tilde{M}_{\mathbb{C}^\times} \rightarrow M_{\mathbb{C}^\times}, \quad (x_i)_{i=1}^{n+2} \mapsto \left( y_i = \prod_{j=1}^{n+2} x_j^{a_{ji}} \right)_{i=1}^{n+2}.$$

The closure of  $\tilde{\mathcal{P}}$  in  $\tilde{M}_{\mathbb{C}}$  is identified with the Milnor fiber

$$(4.3) \quad \check{U} = \{(x_1, \dots, x_{n+2}) \in \mathbb{C}^{n+2} \mid \check{\mathbf{W}}(x_1, \dots, x_{n+2}) + 1 = 0\}$$

of  $\tilde{\mathbf{W}}$ .

We equip  $\check{U}$  with the grading defined by the tensor square  $\Omega_{\check{U}}^{\otimes 2}$  of the holomorphic volume form

$$(4.4) \quad \Omega_{\check{U}} := \text{Res} \frac{dx_1 \wedge \cdots \wedge dx_{n+2}}{\tilde{\mathbf{W}}(x_1, \dots, x_{n+2}) + 1}.$$

If  $n \geq 1$ , then the choice of a grading of  $\check{U}$  is unique because of the simple connectivity of  $\check{U}$ .

The divisors

$$(4.5) \quad E_i = \{(x_1, \dots, x_{n+2}) \in \check{U} \mid x_i = 0\}, \quad i = 1, \dots, n+2$$

are smooth, and one has  $\check{U} \setminus \tilde{\mathcal{P}} = \bigcup_{i=1}^{n+2} E_i$ . If  $n \geq 1$ , then  $E_i$  is non-empty and connected for any  $i = 1, \dots, n+2$ .

Let  $\mathcal{P}^{\text{ua}}$  be the universal abelian cover of  $\mathcal{P}$ , which agrees with the universal cover if  $n \geq 1$ . The inclusion  $\mathcal{P} \hookrightarrow M_{\mathbb{C}^\times}$  induces an isomorphism  $H_1(\mathcal{P}) \xrightarrow{\sim} H_1(M_{\mathbb{C}^\times}) \cong M$ , so that  $M$  is identified with the group  $\text{Deck}(\mathcal{P}^{\text{ua}} \rightarrow \mathcal{P})$  of deck transformations. The group  $\text{Deck}(\mathcal{P}^{\text{ua}} \rightarrow \tilde{\mathcal{P}})$  is naturally identified with  $\tilde{M}$ , so that  $\overline{M} := \text{Deck}(\tilde{\mathcal{P}} \rightarrow \mathcal{P})$  is identified with  $M/\tilde{M}$ .

The Lagrangian immersion  $L^{\text{SS}}$  from an  $n$ -sphere to  $\mathcal{P}$  introduced by Seidel and Sheridan [Sei11, She11] lifts to a Lagrangian immersion  $\tilde{L}^{\text{SS}}$  from the disjoint union of  $|\overline{M}|$  copies of spheres to  $\tilde{\mathcal{P}}$ . Let  $\mathcal{F}_0(U)$  be the full subcategory of  $\mathcal{F}(U)$  split-generated by  $\tilde{L}^{\text{SS}}$ .

Let

$$(4.6) \quad \mathbf{V}(z_0, \dots, z_{n+2}) := \mathbf{W}(z_1, \dots, z_{n+2}) - \prod_{i=0}^{n+2} z_i$$

be a semi-invariant element of  $\mathbf{k}[z_0, \dots, z_{n+2}]$  with respect the natural action of

$$(4.7) \quad K := \left\{ (t_0, \dots, t_{n+2}) \in (\mathbb{G}_m)^{n+2} \left| \prod_{j=1}^{n+2} t_j^{a_{ij}} = \prod_{j=0}^{n+2} t_j \text{ for any } i \in \{1, \dots, n+2\} \right. \right\},$$

and  $\text{mf}_0([\mathbb{A}^{n+2}/K], \mathbf{V})$  be the full subcategory of the dg category  $\text{mf}([\mathbb{A}^{n+2}/K], \mathbf{V})$  of  $K$ -equivariant matrix factorizations of  $\mathbf{V}$  split-generated by the structure sheaf of the origin.

**Proposition 4.2.** *One has an equivalence  $\mathcal{F}_0(U) \simeq \text{mf}_0([\mathbb{A}^{n+2}/K], \mathbf{V})$ .*

*Proof.* Let  $\text{LGr}(\mathcal{P})^{\text{ua}} \rightarrow \text{LGr}(\mathcal{P})$  be the universal abelian cover of  $\text{LGr}(\mathcal{P})$ , whose group of deck transformations can be identified with  $\mathbb{G} := H_1(\text{LGr}(\mathcal{P}))$ . A  $\mathbb{G}$ -grading of a Lagrangian  $L$  in  $\mathcal{P}$  is a lift  $\tilde{s}$  of the tautological section  $s_L: L \rightarrow \text{LGr}(\mathcal{P})$  to  $\text{LGr}(\mathcal{P})^{\text{ua}}$ . The Floer cohomology of  $\mathbb{G}$ -graded Lagrangians is  $\mathbb{G}$ -graded.

The tensor square  $\Omega_{\mathcal{P}}^{\otimes 2}$  of the holomorphic volume form

$$(4.8) \quad \Omega_{\mathcal{P}} := \text{Res} \frac{1}{1 + y_1 + \cdots + y_{n+2}} \frac{dy_1}{y_1} \wedge \cdots \wedge \frac{dy_{n+2}}{y_{n+2}}$$

induces a splitting of the exact sequence

$$(4.9) \quad 0 \rightarrow H_1(\text{LGr}(T_p\mathcal{P})) \rightarrow H_1(\text{LGr}(\mathcal{P})) \rightarrow H_1(\mathcal{P}) \rightarrow 0,$$

where  $\text{LGr}(T\mathcal{P})$  is the Lagrangian Grassmannian bundle of the tangent bundle  $T\mathcal{P}$ , and  $p \in \mathcal{P}$  is an arbitrarily chosen base point. We identify  $\mathbb{G} := H_1(\text{LGr}(\mathcal{P}))$  with  $H_1(\mathcal{P}) \oplus H_1(\text{LGr}(T_p\mathcal{P})) \cong M \oplus \mathbb{Z}$  by this splitting. Similarly, we identify  $\text{Deck}(\text{LGr}(\mathcal{P})^{\text{ua}} \rightarrow \text{LGr}(\tilde{\mathcal{P}}))$  with  $\tilde{M} \oplus \mathbb{Z}$  using the tensor square (of the restriction to  $\tilde{\mathcal{P}}$ ) of  $\Omega_U$ .

The cohomology algebra  $A$  of the endomorphism  $A_\infty$ -algebra  $\mathcal{A}$  of the immersed Lagrangian sphere  $L^{\text{SS}}$  in the  $\mathbb{G}$ -graded Fukaya category of  $\mathcal{P}$  is computed in [She11] as the exterior algebra generated by elements  $\theta_0, \dots, \theta_{n+2}$  of degrees

$$(4.10) \quad \deg \theta_i = \begin{cases} -(\mathbf{e}_1 + \cdots + \mathbf{e}_{n+2}) - 1 & i = 0, \\ 1 + \mathbf{e}_i & i = 1, \dots, n+2. \end{cases}$$

The Kontsevich formality lifts the Hochschild–Kostant–Rosenberg isomorphism to an  $L_\infty$ -quasi-isomorphism

$$(4.11) \quad \Phi_{\text{HKR}}: \text{CC}^\bullet(A) \rightarrow \mathbb{C}[z_0, \dots, z_{n+2}][\theta_0, \dots, \theta_{n+2}]$$

from the Hochschild cochain complex of  $A$  to the graded Lie algebra of polyvector fields on  $\mathbb{C}[z_0, \dots, z_{n+2}]$ . The latter is a formal  $L_\infty$ -algebra, whose underlying graded vector space is the free graded commutative algebra generated by even variables  $z_i$  of degrees

$$(4.12) \quad \deg z_i = -\deg \theta_i + 1$$

and odd variables  $\theta_i$  of degrees (4.10). The non-trivial  $L_\infty$ -operation  $\mathfrak{l}_2$  is identified with the Schouten bracket by sending  $\theta_i$  to  $\frac{\partial}{\partial z_i}$ . As explained in [SS21, Lemma 3.3], the Maurer-Cartan element  $\mu^{\geq 3}$  describing the deformation of  $A$  to  $\mathcal{A}$  is sent to

$$(4.13) \quad \mathbf{V}_0 := -z_0 \cdots z_{n+2}$$

by  $\Phi_{\text{HKR}}$ . As shown in [SS21, Lemma 3.5], this implies that the Hochschild cohomology of  $\mathcal{A}$  is isomorphic to the quotient

$$(4.14) \quad \mathbb{C}[z_0, \dots, z_{n+2}][u_0, \dots, u_{n+2}]/\mathcal{J}$$

of the free graded commutative algebra generated by even variables  $z_i$  of degrees (4.12) and odd variables  $u_i$  of degrees 1 by the ideal

$$(4.15) \quad \mathcal{J} = \left( \prod_{i \notin I} z_i \prod_{i \in I} u_i \right)_{I \subset \{0, \dots, n+2\}}.$$

Let  $\tilde{\mathcal{A}}$  be the endomorphism  $A_\infty$ -algebra of  $\tilde{L}^{\text{SS}}$  in the  $\tilde{\mathbb{G}}$ -graded Fukaya category of  $\tilde{\mathcal{P}}$ . It is obtained from  $\mathcal{A}$  as follows:

- Take a complete set  $\{m_g\}_{g \in \overline{M}} \subset M \subset \mathbb{G} \cong M \oplus \mathbb{Z}$  of representative of  $\overline{M} \cong M/\widetilde{M}$ .
- Take the  $\mathbb{G}$ -graded endomorphism  $A_\infty$ -algebra of the direct sum  $\bigoplus_{g \in \overline{M}} \mathcal{A}(m_g)$  of shifted free modules in  $\text{mod } \mathcal{A}$ .
- Take the  $A_\infty$ -subalgebra consisting of homogeneous elements whose degrees are in  $\tilde{\mathbb{G}}$ .

Let  $\tilde{\mathcal{A}}_R$  be the  $\tilde{\mathbb{G}}$ -graded deformation of  $\tilde{\mathcal{A}}$  over the polynomial ring

$$(4.16) \quad R := \mathbb{C}[r_1, \dots, r_{n+2}]$$

whose  $A_\infty$ -operations are given by counting pseudo-holomorphic disks in  $U$  weighted by intersection numbers with  $E_i$ . The degree

$$(4.17) \quad \deg r_i = \mathbf{f}_i + 2 = \sum_{j=1}^{n+2} a_{ij} \mathbf{e}_j + 2$$

of the variable counting intersection numbers with  $E_i$  is defined by first choosing a small disk in  $U$  intersecting simply and transversely to  $E_i$  and disjoint from all the other  $E_j$ 's, lifting it to  $\text{LGr}(U)$ , and taking the class of its boundary in  $\tilde{\mathbb{G}}$ . Let  $\mathfrak{m} = (r_1, \dots, r_{n+2})$  be the maximal ideal of  $R$  at the origin. The first order deformation class of  $\tilde{\mathcal{A}}_R$  belongs to the  $\overline{M}$ -invariant degree 2 part  $\text{HH}^2(\tilde{\mathcal{A}}, \tilde{\mathcal{A}} \otimes \mathfrak{m}/\mathfrak{m}^2)^{\overline{M}}$  of the  $\tilde{\mathbb{G}}$ -graded Hochschild cohomology of the  $\tilde{\mathcal{A}}$ -bimodule  $\tilde{\mathcal{A}} \otimes \mathfrak{m}/\mathfrak{m}^2$ , which is isomorphic to the degree 2 part  $\text{HH}^2(\mathcal{A}, \mathcal{A} \otimes \mathfrak{m}/\mathfrak{m}^2)$  of the  $\mathbb{G}$ -graded Hochschild cohomology of the  $\mathcal{A}$ -bimodule  $\mathcal{A} \otimes \mathfrak{m}/\mathfrak{m}^2$  by [She15, Remark 2.66].



One has

$$(4.18) \quad \deg \left( \prod_{i=1}^{n+2} r_i^{b_i} \prod_{i=0}^{n+2} z_i^{c_i} \prod_{i \in I} u_i \right)$$

$$(4.19) \quad = \sum_{i=1}^{n+2} b_i \left( \sum_{j=1}^{n+2} a_{ij} \mathbf{e}_j + 2 \right) - \sum_{i=1}^{n+2} c_i \mathbf{e}_i + c_0 (\mathbf{e}_1 + \cdots + \mathbf{e}_{n+2} + 2) + |I|$$

$$(4.20) \quad = \sum_{i=1}^{n+2} \left( \sum_{j=1}^{n+2} a_{ji} b_j - c_i + c_0 \right) \mathbf{e}_i + \sum_{i=1}^{n+2} 2b_i + 2c_0 + |I|.$$

For this to be 2, one needs

$$(4.21) \quad c_i = \sum_{j=1}^{n+2} a_{ji} b_j + c_0$$

for  $i = 1, \dots, n+2$  and

$$(4.22) \quad 2 \sum_{i=1}^{n+2} b_i + 2c_0 + |I| = 2.$$

This is the case if and only if one of  $b_1, \dots, b_n, c_0$ , or  $|I|/2$  is 1 and others are 0. If  $b_j = \delta_{ij}$  for  $i \in \{1, \dots, n+2\}$ , then one has

$$(4.23) \quad c_j = \sum_{k=1}^{n+2} a_{kj} \delta_{ki} = a_{ij}$$

and

$$(4.24) \quad r_i \prod_{j=1}^{n+2} z_j^{c_j} = r_i \prod_{j=1}^{n+2} z_j^{a_{ij}}.$$

This argument also shows  $\mathrm{HH}^2(\mathcal{A}, \mathcal{A} \otimes \mathfrak{m}^i) = 0$  for  $i \geq 2$ .

The first order deformation class of  $\tilde{\mathcal{A}}_R$  is the sum

$$(4.25) \quad \sum_{i=1}^{n+2} r_i \prod_{j=1}^{n+2} z_j^{a_{ij}}$$

of (4.24) for all  $i \in \{1, \dots, n+2\}$ . If  $n > 0$ , this follows from [She20, Lemma 4.22] and [She20, Assumption 5.3].<sup>1</sup> If  $n = 0$ , then one has either

- (1) Brieskorn–Pham:  $\mathbf{W} = \check{\mathbf{W}} = x^p + y^q$ ,
- (2) chain:  $\mathbf{W} = x^p y + y^q$  and  $\check{\mathbf{W}} = x^p + x y^q$ , or
- (3) loop:  $\mathbf{W} = \check{\mathbf{W}} = x^p y + x y^q$ .

In all cases, the first order deformation class can be computed directly, since disk counting in dimension 1 is combinatorial. In the chain case, the divisor in  $\check{U}$  defined by  $x = 0$  is empty and the divisor defined by  $y = 0$  consists of  $p$  points. Correspondingly, one has

$$(4.26) \quad \mathbf{V} = x^p y + y^q - x y z = y^q - x y (z - x^{p-1}),$$

so that the term  $x^p y$  can be absorbed into a coordinate change of  $z$ . Similarly, in the loop case, both of the divisors in  $\check{U}$  defined by  $x = 0$  and  $y = 0$  are empty, and one has

$$(4.27) \quad \mathbf{V} = x^p y + x y^q - x y z = x y (z - x^{p-1} - y^{q-1}),$$

so that both of the terms  $x^p y$  and  $x y^q$  can be absorbed into a coordinate change of  $z$ .

Now  $\tilde{\mathcal{A}}_R$  and hence

$$(4.28) \quad \tilde{\mathcal{A}}_1 := \tilde{\mathcal{A}}_R \otimes_R R / (r_i - 1)_{i=1}^n$$

<sup>1</sup>Although the proof of [She20, Assumption 5.3] is relegated to future work, there is no difficulty in the present setting, where  $U$  is exact.

are determined uniquely up to quasi-isomorphism by [She16, Proposition 6.6], so that

$$(4.29) \quad \mathcal{F}_0(U) \simeq \text{mod } \tilde{\mathcal{A}}_1.$$

Similarly,  $\text{mf}_0(\mathbf{k}[z_0, \dots, z_{n+2}], \mathbf{V}_0)$  admits a  $\mathbb{G}$ -grading by (4.12), and

$$(4.30) \quad \text{mf}_0 \left( R[z_0, \dots, z_{n+2}], \mathbf{V}_R := \mathbf{V}_0 + \sum_{i=1}^{n+2} r_i \prod_{j=1}^{n+2} z_j^{a_{ij}} \right)$$

is a  $\tilde{\mathbb{G}}$ -graded deformation whose first order deformation class is (4.25) by [She15, Proposition 7.1]. It follows that

$$(4.31) \quad \text{mf}_0([\mathbb{A}^{n+2}/K], \mathbf{V}) \simeq \text{mod } \tilde{\mathcal{A}}_1,$$

and Proposition 4.2 is proved.  $\square$

Now we specialize to Brieskorn–Pham singularities where  $a_{ij} = p_i \delta_{ij}$  and  $p_i > 2$  for  $i \in \{1, \dots, n+2\}$ .

**Proposition 4.3.** *One has an equivalence*

$$(4.32) \quad \mathcal{F}_0(\check{U}) \simeq \mathcal{F}(\check{U}).$$

*Proof.* Let

$$(4.33) \quad \varpi: \check{U} \rightarrow \mathbb{C}, \quad (x_k)_{k=1}^{n+2} \mapsto x_{n+2}$$

be the projection to the last coordinate. The fiber

$$(4.34) \quad \varpi^{-1}(x_{n+2}) = \{(x_1, \dots, x_n) \in \mathbb{C}^n \mid x_1^{p_1} + \dots + x_n^{p_n} + (x_{n+2}^{p_{n+2}} + 1) = 0\}$$

is a Milnor fiber of a lower-dimensional Brieskorn–Pham singularity, unless  $x_{n+2}$  belongs to the set

$$(4.35) \quad \text{Critv } \varpi = \left\{ \zeta_k := \exp(2(k+1/2)\pi\sqrt{-1}/p_{n+2}) \right\}_{k=0}^{p_{n+2}-1}$$

of critical values of  $\varpi$ , in which case the fiber has the lower-dimensional Brieskorn–Pham singularity at the origin.

Set

$$(4.36) \quad I := \left\{ \mathbf{i} = (i_1, \dots, i_{n+2}) \in \mathbb{Z}^{n+2} \mid 0 \leq i_k \leq p_k - 2 \text{ for any } 1 \leq k \leq n+2 \right\}.$$

A distinguished basis  $(V_{\mathbf{i}})_{\mathbf{i} \in I}$  of vanishing cycles is described inductively in [FU11] as a fibration of lower-dimensional vanishing cycles above the matching path on the  $x_{n+2}$ -plane connecting  $\zeta_{i_{n+2}}$  and  $\zeta_{i_{n+2}+1}$ .

There is another distinguished basis  $(S_{\mathbf{i}})_{\mathbf{i} \in I}$  of vanishing cycles, which can similarly be constructed inductively as a fibration of lower-dimensional vanishing cycles above the matching path on the  $x_{n+2}$ -plane connecting  $\zeta_0$  and  $\zeta_{i_{n+2}+1}$ . Note that  $S_{\mathbf{i}} = V_{\mathbf{i}}$  for  $\mathbf{i} = \mathbf{0} := (0, \dots, 0)$ . The collection  $(V_{\mathbf{i}})_{\mathbf{i} \in I}$  in the Fukaya–Seidel category of the Brieskorn–Pham polynomial corresponds to simple modules of the tensor product of the path algebras of  $A_{p_k-1}$ -quivers, whereas the collection  $(S_{\mathbf{i}})_{\mathbf{i} \in I}$  corresponds to projective modules.

The collection  $(S_{\mathbf{i}})_{\mathbf{i} \in I}$  of objects in  $\mathcal{F}(\check{U})$  has a Koszul dual collection  $(L_{\mathbf{i}})_{\mathbf{i} \in I}$  of objects in  $\mathcal{W}(\check{U})$ , which are connected components of the inverse image of

$$(4.37) \quad \{(x_1, \dots, x_{n+2}) \in \mathcal{P} \cap \mathbb{R}^{n+2} \mid x_i > 0 \text{ for } 1 \leq i \leq n \text{ and } x_{n+2} < -1\}$$

by the covering map  $\varphi_{\mathbb{C}^\times}: \check{U} \rightarrow \mathcal{P}$ . The Lagrangian  $L_{\mathbf{i}}$  is a Lefschetz thimble for  $\varpi$  over the half line on the  $x_{n+2}$ -planes from  $\zeta_{i_{n+2}+1}$  to infinity associated with the  $(n-1)$ -dimensional vanishing cycle  $V_{\bar{\mathbf{i}}}$  where  $\bar{\mathbf{i}} = (i_k)_{k=1}^n$ .

Since  $L^{\text{SS}}$  intersects (4.37) only at one point [She11, Corollary 2.9], there is an irreducible component  $\tilde{L}_{\mathbf{0}}^{\text{SS}}$  of  $\tilde{L}^{\text{SS}}$  such that  $\tilde{L}_{\mathbf{0}}^{\text{SS}} \cap L_{\mathbf{i}}$  consists of one point if  $\mathbf{i} = \mathbf{0}$ , and is empty otherwise. It follows that  $\tilde{L}_{\mathbf{0}}^{\text{SS}} \simeq S_{\mathbf{0}}$  in  $\mathcal{F}(\check{U})$ . The vanishing cycle  $V_{\mathbf{i}}$  for other  $\mathbf{i}$  is the image of  $V_{\mathbf{0}} = S_{\mathbf{0}}$  by the map  $(x_k)_{k=1}^{n+2} \mapsto (\exp(2i_k\pi\sqrt{-1}/p_k) x_k)_{k=1}^{n+2}$ . Since vanishing cycles generate  $\mathcal{F}(\check{U})$ , Proposition 4.3 is proved.  $\square$

The closed immersion

$$(4.38) \quad \iota: [\mathbf{W}^{-1}(0)/K] \rightarrow [\mathbf{V}^{-1}(0)/K]$$

induces a push-forward

$$(4.39) \quad \iota_*: \text{scoh} [\mathbf{W}^{-1}(0)/K] \rightarrow \text{scoh} [\mathbf{V}^{-1}(0)/K]$$

since the push-forward of the structure sheaf of  $\mathbf{W}^{-1}(0)$  is defined by the non-zero-divisor  $x_0$  and hence has projective dimension one. The adjunction with the pull-back

$$(4.40) \quad \iota^*: \text{scoh} [\mathbf{W}^{-1}(0)/K] \rightarrow \text{scoh} [\mathbf{V}^{-1}(0)/K]$$

follows from that for coherent sheaves. Let

$$(4.41) \quad \mathcal{L}_i := \mathcal{O}_{\mathbb{A}^1 \times \mathbf{0}}(-i_1\chi_1 - \cdots - i_{n+2}\chi_{n+2})$$

be the structure sheaf of the closed subscheme  $\mathbb{A}^1 \times \mathbf{0}$  of  $\mathbf{V}^{-1}(0)$  defined by  $x_1 = \cdots = x_{n+2} = 0$ , twisted by an element of  $\widehat{K} := \text{Hom}(K, \mathbb{G}_m)$  which is generated by

$$(4.42) \quad \chi_i: K \rightarrow \mathbb{G}_m, \quad (\alpha_i)_{i=0}^{n+2} \mapsto \alpha_i^{-1}$$

with relations

$$(4.43) \quad p_1\chi_1 = \cdots = p_{n+2}\chi_{n+2} = \chi_0 + \cdots + \chi_{n+2}.$$

Then the sequence  $(\mathcal{E}_i := \iota^*\mathcal{L}_i)_{i \in I}$  is the full strong exceptional collection given in [FU11]. Let  $(\mathcal{F}_i)_{i \in I}$  be the full exceptional collection right dual to  $(\mathcal{E}_i)_{i \in I}$  so that

$$(4.44) \quad \dim \text{hom}(\mathcal{E}_i, \mathcal{E}_j) = \delta_{i,j},$$

and set

$$(4.45) \quad \mathcal{S}_i := \iota_*\mathcal{F}_i.$$

Then  $(\mathcal{L}_i)_{i \in I}$  and  $(\mathcal{S}_i)_{i \in I}$  are Koszul dual;

$$(4.46) \quad \dim \text{hom}(\mathcal{L}_i, \mathcal{S}_j) \cong \dim \text{hom}(\mathcal{E}_i, \mathcal{F}_j) \simeq \delta_{i,j}.$$

**Proposition 4.4.** *If  $\tilde{\mathbf{w}}$  is not of Calabi–Yau type, then one has an equivalence*

$$(4.47) \quad \text{Fun}^{\text{ex}}(\text{mf}_0([\mathbb{A}^{n+3}/K], \mathbf{V}), \text{perf } \mathbf{k}) \simeq \text{mf}([\mathbb{A}^{n+3}/K], \mathbf{V})$$

of  $\infty$ -categories.

*Proof.* This follows from (4.46) and the fact that the right orthogonal to  $(\mathcal{S}_i)_{i \in I}$  is zero since  $\mathbb{G}_m \subset K$  is a dilating action on the critical locus  $\mathbb{A}^1 \times \mathbf{0}$  of  $\mathbf{V}$  with the origin as the unique fixed point.  $\square$

Proposition 4.4 combined with [LU22, Theorem 6.11] proves [LU22, Conjecture 1.4] for non-log Calabi–Yau Brieskorn–Pham singularity.

## 5. RABINOWITZ FLOER HOMOLOGY FROM MIRROR SYMMETRY

5.1. We use the same notations as in Section 3. We regard  $x_0$  as a section of the line bundle  $\mathcal{L}$  on  $U$  associated with the character  $\chi_0$  of  $K$ , which in turn gives a natural transformation  $s$  from the autoequivalence  $\mathcal{L}^\vee \otimes (-) \simeq (-\chi_0)$  on  $\text{mf}([\mathbb{A}^{n+3}/K], \mathbf{W} - x_0 \cdots x_{n+2}) \simeq \text{scoh } U$  to the identity functor. The localization of  $\text{scoh } U$  along  $s$  gives  $\text{scoh } D \simeq \text{scoh } U \setminus E$  since  $E$  is defined by  $x_0 = 0$  (cf. [Sei08, Section 1]).

Let  $(d_1, \dots, d_{n+2}, h)$  be the sequence of positive integers such that  $\gcd(d_1, \dots, d_{n+2}, h) = 1$  and

$$(5.1) \quad \mathbf{W}(t^{d_1}x_1, \dots, t^{d_{n+2}}x_{n+2}) = t^h\mathbf{W}(x_1, \dots, x_{n+2}).$$

Then  $h$  is the minimal positive integer such that  $x_0^h$  is invariant under the action of  $\ker \chi \subset K$ , and one has

$$(5.2) \quad (\mathcal{L}^\vee)^{\otimes h} \otimes (-) \simeq (-h\chi_0) \simeq (-d_0\chi) \simeq [-2d_0]$$

where

$$(5.3) \quad d_0 := h - d_1 - \cdots - d_{n+2}.$$

One can regard  $s^h$  as an element of  $\mathrm{HH}^{2d_0}(\mathrm{scoh} U)$ , so that  $\mathrm{scoh} U$  is linear over  $\mathbf{k}[s^h]$ . The localization of  $\mathrm{scoh} U$  along  $s$  is equivalent to that along  $s^h$ ;

$$(5.4) \quad \mathrm{scoh} D \simeq \mathrm{scoh} U \otimes_{\mathbf{k}[s^h]} \mathbf{k}[s^h, s^{-h}].$$

Assume Conjecture 3.2, so that  $\mathrm{scoh} D \simeq \mathcal{R}(\check{U})$  by (3.19). This implies

$$(5.5) \quad \mathrm{RFH}^*(\check{U}) \simeq \mathrm{HH}^*(\mathcal{R}(\check{U}))$$

$$(5.6) \quad \simeq \mathrm{HH}^*(\mathrm{scoh} U \otimes_{\mathbf{k}[s^h]} \mathbf{k}[s^h, s^{-h}])$$

$$(5.7) \quad \simeq \mathrm{HH}^*(\mathrm{scoh} U) \otimes_{\mathbf{k}[s^h]} \mathbf{k}[s^h, s^{-h}].$$

5.2. Let  $\check{\mathbf{W}}$  be the transpose of  $\mathbf{W}$  and  $(\check{d}_1, \dots, \check{d}_{n+2}, \check{h})$  be the sequence of positive integers such that  $\mathrm{gcd}(\check{d}_1, \dots, \check{d}_{n+2}, \check{h}) = 1$  and

$$(5.8) \quad \check{\mathbf{W}}(t^{\check{d}_1}x_1, \dots, t^{\check{d}_{n+2}}x_{n+2}) = t^{\check{h}}\check{\mathbf{W}}(x_1, \dots, x_{n+2}).$$

Then the link

$$(5.9) \quad C := \{(x_1, \dots, x_{n+2}) \in \check{\mathbf{W}}^{-1}(0) \mid |x_1|^2 + \dots + |x_{n+2}|^2 = 1\}$$

of the singularity of  $\check{\mathbf{W}}^{-1}(0)$  at the origin has an  $S^1$ -action defined by

$$(5.10) \quad S^1 \ni t: (x_1, \dots, x_{n+2}) \mapsto (t^{\check{d}_1}x_1, \dots, t^{\check{d}_{n+2}}x_{n+2}),$$

which lifts to an  $S^1$ -action

$$(5.11) \quad S^1 \ni t: (x_0, \dots, x_{n+2}) \mapsto (t^{\check{h}}x_0, t^{\check{d}_1}x_1, \dots, t^{\check{d}_{n+2}}x_{n+2})$$

on the total space of the family

$$(5.12) \quad \varphi: \check{\mathcal{U}} := \{(x_0, \dots, x_{n+2}) \in \mathbb{C}^{n+3} \mid \check{\mathbf{W}}(x_1, \dots, x_{n+2}) = x_0\} \rightarrow \mathbb{C}^1, \quad (x_0, \dots, x_{n+2}) \mapsto x_0.$$

Let  $\mu$  be the endofunctor of  $\mathcal{W}(\check{U})$  defined as the clockwise monodromy of the family (5.12) around the origin, which is isomorphic to the composite of inverse spherical twists along a distinguished basis of vanishing cycles of  $\check{\mathbf{W}}$ . It is shown in [Sei00, Section 4.c] that

$$(5.13) \quad \mu^{\check{h}} \simeq [-2\check{d}_0]$$

where

$$(5.14) \quad \check{d}_0 := \check{h} - \check{d}_1 - \dots - \check{d}_{n+2}.$$

The wrapped Fukaya category of the singular hypersurface  $\check{\mathbf{W}}^{-1}(0)$  in the sense of Auroux (cf. [Jef22, Definition 1]), defined as the localization of  $\mathcal{W}(\check{U})$  along the natural transformation  $\check{s}: \mu \rightarrow \mathrm{id}$  first introduced in [Sei09], is equivalent to the quotient of  $\mathcal{W}(\check{U})$  by the split-closure of the essential image of the cap functor [Jef22, Corollary 1]. Since  $\mathcal{F}(\check{U})$  is split-generated by vanishing cycles, which are in the essential image of the cap functor, the wrapped Fukaya category of the singular hypersurface  $\check{\mathbf{W}}^{-1}(0)$  is equivalent to the stable Fukaya category, which in turn is equivalent to the Rabinowitz Fukaya category  $\mathcal{R}(\check{U})$ .

5.3. Set

$$(5.15) \quad V := \mathbf{k}x_0 \oplus \mathbf{k}x_1 \oplus \dots \oplus \mathbf{k}x_{n+2}.$$

For  $\gamma \in K$ , let  $V_\gamma$  be the subspace of  $\gamma$ -invariant elements in  $V$ ,  $S_\gamma$  be the symmetric algebra of  $V_\gamma$ ,  $\mathbf{V}_\gamma$  be the restriction of  $\mathbf{V}$  to  $\mathrm{Spec} S_\gamma$ , and  $N_\gamma$  be the  $K$ -stable complement of  $V_\gamma$  in  $V$  so that

$V \cong V_\gamma \oplus N_\gamma$  as a  $K$ -module. Then [Dyc11, CT13, Seg13, BFK14] (cf. also [LU22, Theorem 3.1]) shows that  $\mathrm{HH}^t(\mathrm{mf}([\mathbb{A}^{n+2}/K], \mathbf{V}))$  is isomorphic to

$$(5.16) \quad \bigoplus_{\substack{\gamma \in \ker \chi, l \geq 0 \\ t - \dim N_\gamma = 2u}} (H^{-2l}(d\mathbf{V}_\gamma) \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee)_{(u+l)\chi} \\ \oplus \bigoplus_{\substack{\gamma \in \ker \chi, l \geq 0 \\ t - \dim N_\gamma = 2u+1}} (H^{-2l-1}(d\mathbf{V}_\gamma) \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee)_{(u+l+1)\chi}.$$

Here  $H^i(d\mathbf{V}_\gamma)$  is the  $i$ -th cohomology of the Koszul complex

$$(5.17) \quad C^*(d\mathbf{V}_\gamma) := \{ \cdots \rightarrow \Lambda^2 V_\gamma^\vee \otimes S_\gamma(-2\chi) \rightarrow V_\gamma^\vee \otimes S_\gamma(-\chi) \rightarrow S_\gamma \},$$

where the rightmost term  $S_\gamma$  sits in cohomological degree 0, and the differential is the contraction with

$$(5.18) \quad d\mathbf{V}_\gamma \in (V_\gamma \otimes S_\gamma)_\chi.$$

5.4. As an example, consider the case  $\mathbf{W} = x_1^2 + \cdots + x_{n+2}^2$ , where  $\check{U} \cong T^*S^{n+1}$ ,  $h = 2$ ,  $d_0 = -n$ , and  $\mathrm{HH}^*(\mathrm{mf}([\mathbb{A}^5/K], \mathbf{W})) \otimes_{\mathbf{k}[s^h]} \mathbf{k}[s^h, s^{-h}]$  has a basis consisting of

- $x_0^{2m}$  for  $m \in \mathbb{Z}$  and  $\gamma = (1, \dots, 1)$  of degree  $-2mn$ ,
- $x_0^{2m+1} \otimes x_0^\vee$  for  $m \in \mathbb{Z}$  and  $\gamma = (1, \dots, 1)$  of degree  $-2mn + 1$

and, if  $n$  is even, in addition to the above,

- $x_0^{2m} \otimes x_1^\vee \otimes \cdots \otimes x_{n+2}^\vee$  for  $m \in \mathbb{Z}$  and  $\gamma = (1, -1, -1, \dots, -1)$  of degree  $(2m+1)n$ ,
- $x_0^{2m+1} \otimes x_0^\vee \otimes \cdots \otimes x_{n+2}^\vee$  for  $m \in \mathbb{Z}$  and  $\gamma = (1, -1, -1, \dots, -1)$  for  $m \in \mathbb{Z}$  of degree  $(2m+1)n+1$ .

5.5. Let  $U_n$  be the Liouville domain  $\mathbb{C}^\times \cong T^*S^1$  equipped with the grading determined by the quadratic differential  $x^n(d \log x)^{\otimes 2}$ . Then  $\mathcal{W}(U_n)$  is generated by the cotangent fiber, whose endomorphism  $A_\infty$ -algebra is isomorphic to the free algebra  $\mathbf{k}\langle u, u^{-1} \rangle$  generated by an element  $u$  of degree  $n$  and its inverse  $u^{-1}$ , so that

$$(5.19) \quad \mathcal{W}(U_n) \simeq \mathrm{mod} \mathbf{k}\langle u, u^{-1} \rangle$$

$$(5.20) \quad \simeq \mathcal{R}(T^*S^{n+1}),$$

and hence

$$(5.21) \quad \mathrm{SH}^*(U_n) \simeq \mathrm{HH}^*(\mathbf{k}\langle u, u^{-1} \rangle)$$

$$(5.22) \quad \simeq \mathrm{RFH}(T^*S^{n+1}).$$

Reeb orbits in the contact boundary of  $U_n$  with winding number  $w$  come in a family parametrized by  $S^1$ , and a Bott–Morse model of the symplectic cohomology gives two generators  $p_w$  and  $q_w$  of degrees  $nw$  and  $nw + 1$  in such a way that the Floer differential is given by

$$(5.23) \quad dp_w = (1 - (-1)^{nw})q_w$$

(see [BO09, Propostion 3.9]). It follows that

- if  $n$  is even, then  $p_w$  and  $q_w$  for all  $w \in \mathbb{Z}$  survives,
- if  $n$  is odd, then  $p_w$  and  $q_w$  for odd  $w$  annihilates each other, and only  $p_w$  and  $q_w$  for even  $w$  survives,

and one can identify

$$(5.24) \quad p_{2m} = x_0^{2m},$$

$$(5.25) \quad q_{2m} = x_0^{2m+1} \otimes x_0^\vee,$$

$$(5.26) \quad p_{2m+1} = x_0^m \otimes x_1^\vee \otimes \cdots \otimes x_{n+2}^\vee,$$

$$(5.27) \quad q_{2m+1} = x_0^{2m+1} x_0^\vee \otimes x_1^\vee \otimes \cdots \otimes x_{n+2}^\vee.$$

5.6. One can also compute  $\mathrm{HH}^*(\mathbf{k}\langle u, u^{-1} \rangle)$  in a purely algebraic way. The enveloping algebra  $R^{\mathrm{op}} \otimes R$  of  $R := \mathbf{k}\langle u, u^{-1} \rangle$  is isomorphic to  $E = \mathbf{k}\langle \lambda^{\pm 1}, \rho^{\pm 1} \rangle / (\lambda\rho - (-1)^n \rho\lambda)$ . The diagonal bimodule  $\Delta$  regarded as a right  $E$ -module is the  $\mathbf{k}$ -vector space  $\bigoplus_{i=-\infty}^{\infty} \mathbf{k}u^i$  equipped with the action

$$(5.28) \quad u^i \cdot \lambda = (-1)^{in} u^{i+1},$$

$$(5.29) \quad u^i \cdot \rho = u^{i+1}.$$

One has

$$(5.30) \quad \Delta \simeq \left\{ E[-n] \xrightarrow{(\lambda-\rho)\cdot} E \right\},$$

where the right terms sits in degree zero, so that the Hochschild complex of  $R$  is given by

$$(5.31) \quad \mathrm{hom}_E(\Delta, \Delta) \simeq \mathrm{hom}_E \left( \left\{ E[-n] \xrightarrow{(\lambda-\rho)\cdot} E \right\}, \Delta \right)$$

$$(5.32) \quad \simeq \Delta \otimes_E \left\{ E \xrightarrow{(\lambda-\rho)\cdot} E[n] \right\}$$

$$(5.33) \quad \simeq \left\{ \Delta \xrightarrow{(\lambda-\rho)\cdot} \Delta[n] \right\},$$

where the left term sits in degree zero. Since

$$(5.34) \quad u^i \cdot (\lambda - \rho) = \begin{cases} 0 & i \text{ is even,} \\ (1 - (-1)^{in})u^{i+1} & i \text{ is odd,} \end{cases}$$

the complex (5.33) is identical to the Floer complex appearing in Section 5.5.

5.7. As another example, consider the case

$$(5.35) \quad \mathbf{W}(x_1, x_2, x_3, x_4) = x_1^k + x_2^{m-k} + x_3^2 + x_4^2.$$

The hypersurface  $\mathbf{W}^{-1}(0)$  has an isolated cDV singularity at the origin.

5.8. Recall from [Rei83, Theorem 1.1] that a 3-fold singularity is terminal of index 1 if and only if it is an isolated cDV singularity. It follows from [McL16, Theorem 1.1] that the link of a terminal singularity is index-positive in the sense of [CO18, Section 9.5], so that the Rabinowitz Floer cohomology is an invariant of the link (i.e., does not depend on the symplectic filling) (see [CO18, Proposition 9.17]).

5.9. One can find a formal change of coordinates transforming  $\mathbf{V}(x_0, x_1, x_2, x_3, x_4)$  to  $\mathbf{W}(x_1, x_2, x_3, x_4)$  to show  $\mathrm{mf}([\mathbb{A}^5/K], \mathbf{V}) \simeq \mathrm{mf}([\mathbb{A}^5/K], \mathbf{W})$  just as in Section 3. Explicit computations of  $\dim \mathrm{HH}^t(\mathrm{mf}([\mathbb{A}^5/K], \mathbf{W}))$  are discussed for simple singularities in [LU21, Section 5], for more general cases in [EL23, Section 3.1], and for all of (5.35) in [APZ, Theorem C].

5.10. One has  $\mathrm{HH}^t(\mathrm{mf}([\mathbb{A}^5/K], \mathbf{W})) = 0$  for  $t > 3$ , and a basis of  $\mathrm{HH}^3(\mathrm{mf}([\mathbb{A}^5/K], \mathbf{W}))$  consists of

$$(5.36) \quad x_0^\vee x_1^\vee x_2^\vee x_3^\vee x_4^\vee \in H^0(C^*(d\mathbf{W}_\gamma)) \otimes \Lambda^5 N_\gamma^\vee \cong \Lambda^5 N_\gamma^\vee$$

in the direct summand of (5.16) such that  $V_\gamma = 0$  and

$$(5.37) \quad x_0^\vee x_1^\vee x_2^\vee x_3^\vee x_4^\vee \in H^{-1}(C^*(d\mathbf{W}_\gamma)) \otimes \Lambda^4 N_\gamma^\vee \cong \mathbf{k}[x_0] \otimes (\mathbf{k}x_0)^\vee \otimes \Lambda^4 N_\gamma^\vee$$

in the direct summand of (5.16) such that  $V_\gamma = \mathbf{k}x_0$ . It follows that  $\dim \mathrm{HH}^3(\mathrm{mf}([\mathbb{A}^5/K], \mathbf{W}))$  is equal to the number  $(m - k - 1)(k - 1)$  of

$$(5.38) \quad \gamma \in \ker \chi = \{(t_0, t_1, t_2, t_3, t_4) \in (\mathbf{k}^\times)^5 \mid t_1^k = t_2^{m-k} = t_3^2 = t_4^2 = t_0 t_1 t_2 t_3 t_4 = 1\}$$

such that  $t_i \neq 1$  for  $i = 1, 2, 3, 4$ , which in turn is equal to the Milnor number of the singularity defined by (5.35). One also has  $\mathrm{HH}^2(\mathrm{mf}([\mathbb{A}^5/K], \mathbf{W})) = 0$  and  $\dim \mathrm{HH}^{2i}(\mathrm{mf}([\mathbb{A}^5/K], \mathbf{W})) = \dim \mathrm{HH}^{2i+1}(\mathrm{mf}([\mathbb{A}^5/K], \mathbf{W}))$  for all  $i \leq 0$ . A basis of  $\mathrm{HH}^{-2i}(\mathrm{mf}([\mathbb{A}^5/K], \mathbf{W}))$  for  $i \in \mathbb{N}$  can be divided into two kinds.

5.11. Let  $m_1: \mathbb{N} \rightarrow \mathbb{N}$  and  $b_1: \mathbb{N} \rightarrow \{0, \dots, k-1\}$  (resp.  $m_2: \mathbb{N} \rightarrow \mathbb{N}$  and  $b_2: \mathbb{N} \rightarrow \{0, \dots, k-m-1\}$ ) be the quotient and the remainder by  $k$  (resp.  $m-k$ ), so that

$$(5.39) \quad b_0 = b_1(b_0) + km_1(b_0) = b_2(b_0) + (m-k)m_2(b_0)$$

for any  $b_0 \in \mathbb{N}$ . A basis of the first kind in cohomological degree  $-2i$  for  $i \in \mathbb{N}$  is parametrized by the set

$$(5.40) \quad \mathbb{I}_{k,m-k}^i := \{b_0 \in \mathbb{N} \mid b_1(b_0) \neq k-1, b_2(b_0) \neq m-k-1, m_1(b_0) + m_2(b_0) = i\}$$

as

$$(5.41) \quad \begin{cases} x_0^{b_0} x_1^{b_1(b_0)} x_2^{b_2(b_0)} & b_0 \text{ is even and } \gamma = (1, 1, 1, 1, 1), \\ x_0^{b_0} x_1^{b_1(b_0)} x_2^{b_2(b_0)} x_3^\vee x_4^\vee & b_0 \text{ is odd and } \gamma = (1, 1, 1, -1, -1). \end{cases}$$

5.12. A basis of the second kind in cohomological degree  $-2i$  is parametrized by the product set of

$$(5.42) \quad \mathbb{II}_{k,m-k}^i := \{b_0 \in \mathbb{N} \mid \exists n_1, n_2 \in \mathbb{N}, n_1 + n_2 - 1 = i \text{ and } b_0 = -1 + kn_1 = -1 + (m-k)n_2\}$$

and

$$(5.43) \quad (\boldsymbol{\mu}_k \times \boldsymbol{\mu}_{m-k}) \setminus \{1\} = \{\xi \in \mathbf{k} \mid \xi \neq 1 \text{ and } \xi^k = \xi^{m-k} = 1\}$$

as

$$(5.44) \quad \begin{cases} x_0^{b_0} x_1^\vee x_2^\vee & b_0 \text{ is even and } \gamma = (1, \xi, \xi^{-1}, 1, 1), \\ x_0^{b_0} x_1^\vee x_2^\vee x_3^\vee x_4^\vee & b_0 \text{ is odd and } \gamma = (1, \xi, \xi^{-1}, -1, -1). \end{cases}$$

5.13. Table 1 shows  $\dim \mathrm{HH}^t(\mathrm{mf}([\mathbb{A}^5/K], \mathbf{W}))$  for  $3 \geq t \geq -15$  for small  $(m, k)$  to give an idea about the outcome of this computation.

$(m, k)$	3	2	1	0	-1	-2	-3	-4	-5	-6	-7	-8	-9	-10	-11	-12	-13	-14	-15
(4, 2)	1	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
(5, 2)	2	0	1	1	0	0	0	0	1	1	0	0	1	1	0	0	0	0	1
(6, 2)	3	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
(7, 2)	4	0	1	1	1	1	0	0	0	0	1	1	1	1	0	0	1	1	1
(6, 3)	4	0	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
(7, 3)	6	0	2	2	0	0	1	1	1	1	0	0	2	2	0	0	2	2	0
(8, 3)	8	0	2	2	1	1	0	0	2	2	0	0	1	1	2	2	0	0	2
(8, 4)	9	0	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3

TABLE 1. Examples of  $\dim \mathrm{HH}^t(\mathrm{mf}([\mathbb{A}^5/K], \mathbf{W}))$

5.14. Define

$$(5.45) \quad h_{k,m-k}^i := \dim \mathrm{HH}^{-2i}(\mathrm{mf}([\mathbb{A}^5/K], \mathbf{W})).$$

Set  $\ell = \gcd(k, m-k)$  and write  $(k, m-k) = (e\ell, f\ell)$ , so that  $\gcd(e, f) = 1$ .

**Proposition 5.1.** *One has*

$$(5.46) \quad h_{e\ell, f\ell}^i = \ell h_{e,f}^i + \ell - 1.$$

*Proof.* If  $i \not\equiv -1 \pmod{e+f}$ , then one has  $\mathbb{II}_{e\ell, f\ell}^i = \emptyset$  so that

$$(5.47) \quad h_{e\ell, f\ell}^i = |\mathbb{I}_{e\ell, f\ell}^i|.$$

If we write

$$(5.48) \quad \mathbb{I}_{e,f}^i = \{b_0^{\min}, b_0^{\min} + 1, \dots, b_0^{\max}\},$$

then one has either

- $b_1(b_0^{\min}) = 0$  and  $b_2(b_0^{\max}) = f - 2$ , or

- $b_2(b_0^{\min}) = 0$  and  $b_1(b_0^{\max}) = e - 2$ .

In both cases, one has

$$(5.49) \quad I_{\ell e, \ell f}^i = \{\ell b_0^{\min}, b_0^{\min} + 1, \dots, \ell b_0^{\max} + 2\ell - 2\},$$

so that

$$(5.50) \quad |I_{\ell e, \ell f}^i| = (\ell b_0^{\max} + 2\ell - 2) - \ell b_0^{\min} + 1$$

$$(5.51) \quad = \ell(b_0^{\max} - b_0^{\min} + 1) + \ell - 1$$

$$(5.52) \quad = \ell |I_{e, f}^i| + \ell - 1.$$

If  $i \equiv -1 \pmod{e + f}$ , then one has  $I_{\ell e, \ell f}^i = \emptyset$ ,  $|I_{\ell e, \ell f}^i| = 1$ , and

$$(5.53) \quad h_{\ell e, \ell f}^i = |(\mu_{\ell e} \cap \mu_{\ell f}) \setminus \{1\}| = \ell - 1,$$

which is equal to  $\ell h_{e, f}^i + \ell - 1$  since  $h_{e, f}^i = 0$ . □

5.15. The following conjecture is proposed in [EL23] (see also [Pet]):

**Conjecture 5.2** ([EL23, Conjecture 1.4]). *A compound Du Val singularity admits a small resolution if and only if the dimension of the symplectic cohomology of its Milnor fiber is constant in every negative cohomological degree. Furthermore, if this is the case, then this dimension is equal to the number of irreducible components of the exceptional locus of a small resolution.*

The following refinement of Conjecture 5.2 was also proposed by the authors of [EL23] based on unpublished calculations:

**Conjecture 5.3.** *Let  $Y \rightarrow X$  be a small  $\mathbb{Q}$ -factorialization of a compound Du Val singularity  $P \in X$ . Let further  $\ell$  be the number of irreducible components of the exceptional locus,  $\check{U}$  be the Milnor fiber of  $P \in X$ , and  $\check{U}_1, \dots, \check{U}_r$  be the Milnor fibers of the resulting  $\mathbb{Q}$ -factorial singularities  $Q_1, \dots, Q_r \in Y$ . Then one has*

$$(5.54) \quad \dim \mathrm{SH}^i(\check{U}) = \sum_{j=1}^r \dim \mathrm{SH}^i(\check{U}_j) + \ell$$

for any  $i < 0$ .

5.16. The blow-up of

$$(5.55) \quad \mathrm{Spec} \mathbf{k}[x, y, z, w]/(xy - f(z, w)g(z, w))$$

along the ideal  $(x, f(z, w))$  is defined by  $xv = f(z, w)u$  and  $yu = g(z, w)v$  inside  $\mathbb{A}_{x, y, z, w}^4 \times \mathbb{P}_{u, v}^1$ , which is the union of

$$(5.56) \quad \mathrm{Spec} \mathbf{k}[x, z, w, v]/(xv - f(z, w))$$

and

$$(5.57) \quad \mathrm{Spec} \mathbf{k}[y, z, w, u]/(yu - g(z, w)).$$

5.17. The 3-fold

$$(5.58) \quad \mathrm{Spec} \mathbf{k}[x, y, z, w]/(x^2 + y^2 + z^{\ell} + w^{\ell}) \cong \mathrm{Spec} \mathbf{k}[x, y, z, w] \Big/ \left( xy - \prod_{i=0}^{\ell-1} (z^e - \zeta_{\ell}^i w^f) \right)$$

where  $\zeta_{\ell} := \exp(2\pi\sqrt{-1}/\ell)$  can be blown-up  $\ell - 1$  times to obtain a chain of  $\ell - 1$  exceptional  $\mathbb{P}^1$ 's and  $\ell$  copies of  $\mathbb{Q}$ -factorial compound Du Val singularities isomorphic to  $x^2 + y^2 + z^e + w^f$ . Proposition 5.1 shows that Conjecture 5.3 holds for the singularity defined by (5.35).



5.18. For  $\mathbf{W}$  given by (5.35), one has

$$(5.59) \quad (h, d_0) = \begin{cases} (ef\ell/2, -e - f) & \ell \text{ is even,} \\ (ef\ell, -2e - 2f) & \ell \text{ is odd.} \end{cases}$$

The isomorphism  $\mathrm{HH}^*(\mathrm{scoh} D) \simeq \mathrm{HH}^*(\mathrm{scoh} U) \otimes_{\mathbf{k}[s^h]} \mathbf{k}[s^h, s^{-h}]$  and the explicit description of  $\mathrm{HH}^*(\mathrm{scoh} U) \cong \mathrm{HH}^*(\mathrm{mf}([\mathbb{A}^5/K], \mathbf{W}))$  above imply  $\mathrm{HH}^i(\mathrm{scoh} D) \cong \mathrm{HH}^i(\mathrm{scoh} U)$  for all  $i \leq 0$  and that the isomorphisms  $s^h: \mathrm{HH}^i(\mathrm{scoh} U) \xrightarrow{\sim} \mathrm{HH}^{i+2d_0}(\mathrm{scoh} U)$  for  $i \leq 0$  extend to isomorphisms  $s^h: \mathrm{HH}^i(\mathrm{scoh} D) \xrightarrow{\sim} \mathrm{HH}^{i+2d_0}(\mathrm{scoh} D)$  for all  $i \in \mathbb{Z}$ . It follows that the symplectic cohomology and the Rabinowitz Floer cohomology of the mirror  $\check{U}$  satisfies  $\mathrm{RFH}^i(\check{U}) \cong \mathrm{SH}^i(\check{U})$  for all  $i \leq 0$  and  $\mathrm{RFH}^i(\check{U}) \cong \mathrm{RFH}^{i+2d_0}(\check{U})$  for all  $i \in \mathbb{Z}$ .

5.19. The non-vanishing of the Rabinowitz Floer homology implies the non-displaceability of the contact boundary of  $\check{U}$  by [CF09, Theorem 1.2].

## REFERENCES

- [APZ] Nikolas Adaloglou, Federica Pasquotto, and Aline Zanardini, *Symplectic cohomology of  $cA_n$  singularities*, arXiv:2404.17301. 14
- [AR87] Maurice Auslander and Idun Reiten, *Almost split sequences for  $\mathbf{Z}$ -graded rings*, Singularities, representation of algebras, and vector bundles (Lambrecht, 1985), Lecture Notes in Math., vol. 1273, Springer, Berlin, 1987, pp. 232–243. MR 915178 (89b:13031) 3
- [BFK14] Matthew Ballard, David Favero, and Ludmil Katzarkov, *A category of kernels for equivariant factorizations and its implications for Hodge theory*, Publ. Math. Inst. Hautes Études Sci. **120** (2014), 1–111. MR 3270588 13
- [BH93] Per Berglund and Tristan Hübsch, *A generalized construction of mirror manifolds*, Nuclear Phys. B **393** (1993), no. 1-2, 377–391. MR MR1214325 (94k:14031) 2
- [BJK] Hanwool Bae, Wonbo Jeong Jeong, and Jongmyeong Kim, *Cluster categories from Fukaya categories*, arXiv:2209.09442. 3
- [BKM16] Karin Baur, Alastair D. King, and Bethany R. Marsh, *Dimer models and cluster categories of Grassmannians*, Proc. Lond. Math. Soc. (3) **113** (2016), no. 2, 213–260. MR 3534972 1
- [BO09] Frédéric Bourgeois and Alexandru Oancea, *Symplectic homology, autonomous Hamiltonians, and Morse-Bott moduli spaces*, Duke Math. J. **146** (2009), no. 1, 71–174. MR 2475400 13
- [Buc87] Ragnar-Olaf Buchweitz, *Maximal Cohen-Macaulay modules and Tate-cohomology over Gorenstein rings*, Available from <https://tspace.library.utoronto.ca/handle/1807/16682>, 1987. 1
- [CF09] Kai Cieliebak and Urs Adrian Frauenfelder, *A Floer homology for exact contact embeddings*, Pacific J. Math. **239** (2009), no. 2, 251–316. MR 2461235 17
- [CO18] Kai Cieliebak and Alexandru Oancea, *Symplectic homology and the Eilenberg-Steenrod axioms*, Algebr. Geom. Topol. **18** (2018), no. 4, 1953–2130, Appendix written jointly with Peter Albers. MR 3797062 14
- [CT13] Andrei Căldăraru and Junwu Tu, *Curved  $A_\infty$  algebras and Landau-Ginzburg models*, New York J. Math. **19** (2013), 305–342. MR 3084707 13
- [Dyc11] Tobias Dyckerhoff, *Compact generators in categories of matrix factorizations*, Duke Math. J. **159** (2011), no. 2, 223–274. MR 2824483 (2012h:18014) 13
- [Eis80] David Eisenbud, *Homological algebra on a complete intersection, with an application to group representations*, Trans. Amer. Math. Soc. **260** (1980), no. 1, 35–64. MR MR570778 (82d:13013) 1
- [EL23] Jonathan David Evans and Yanki Lekili, *Symplectic cohomology of compound Du Val singularities*, Ann. H. Lebesgue **6** (2023), 727–765. MR 4648096 14, 16
- [ET11] Wolfgang Ebeling and Atsushi Takahashi, *Strange duality of weighted homogeneous polynomials*, Compos. Math. **147** (2011), no. 5, 1413–1433. MR 2834726 3
- [FKK23] David Favero, Daniel Kaplan, and Tyler L. Kelly, *Exceptional collections for mirrors of invertible polynomials*, Math. Z. **304** (2023), no. 2, Paper No. 32, 16. MR 4591879 3
- [FU11] Masahiro Futaki and Kazushi Ueda, *Homological mirror symmetry for Brieskorn-Pham singularities*, Selecta Math. (N.S.) **17** (2011), no. 2, 435–452. MR 2803848 (2012e:14083) 1, 3, 10, 11
- [FU13] ———, *Homological mirror symmetry for singularities of type D*, Math. Z. **273** (2013), no. 3-4, 633–652. MR 3030671 3
- [Gam24] Benjamin Gammage, *Mirror symmetry for Berglund-Hübsch Milnor fibers*, Adv. Math. **443** (2024), Paper No. 109563, 45. MR 4713718 5
- [GGV] Sheel Ganatra, Yuan Gao, and Sara Venkatesh, *Rabinowitz Fukaya categories and the categorical formal punctured neighborhood of infinity*, arXiv:2212.14863. 2
- [Hab22] Matthew Habermann, *Homological mirror symmetry for invertible polynomials in two variables*, Quantum Topol. **13** (2022), no. 2, 207–253. MR 4404801 5

- [HLW23] Yang Han, Xin Liu, and Kai Wang, *Exact Hochschild extensions and deformed Calabi-Yau completions*, *Comm. Algebra* **51** (2023), no. 2, 757–778. MR 4532824 4
- [HO23] Yuki Hirano and Genki Ouchi, *Derived factorization categories of non-Thom-Sebastiani-type sums of potentials*, *Proc. Lond. Math. Soc.* (3) **126** (2023), no. 1, 1–75. MR 4535018 3
- [HS20] Matthew Habermann and Jack Smith, *Homological Berglund-Hübsch mirror symmetry for curve singularities*, *J. Symplectic Geom.* **18** (2020), no. 6, 1515–1574. MR 4325377 3
- [IT13] Osamu Iyama and Ryo Takahashi, *Tilting and cluster tilting for quotient singularities*, *Math. Ann.* **356** (2013), no. 3, 1065–1105. MR 3063907 4
- [Iya18] Osamu Iyama, *Tilting Cohen-Macaulay representations*, *Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. II. Invited lectures*, World Sci. Publ., Hackensack, NJ, 2018, pp. 125–162. MR 3966760 3
- [Jef22] Maxim Jeffs, *Mirror symmetry and Fukaya categories of singular hypersurfaces*, *Adv. Math.* **397** (2022), Paper No. 108116, 36. MR 4377932 12
- [JKS16] Bernt Tore Jensen, Alastair D. King, and Xiuping Su, *A categorification of Grassmannian cluster algebras*, *Proc. Lond. Math. Soc.* (3) **113** (2016), no. 2, 185–212. MR 3534971 1
- [Kel05] Bernhard Keller, *On triangulated orbit categories*, *Doc. Math.* **10** (2005), 551–581. MR 2184464 (2007c:18006) 4
- [Kel11] ———, *Deformed Calabi-Yau completions*, *J. Reine Angew. Math.* **654** (2011), 125–180, With an appendix by Michel Van den Bergh. MR 2795754 4
- [Knö87] Horst Knörrer, *Cohen-Macaulay modules on hypersurface singularities. I*, *Invent. Math.* **88** (1987), no. 1, 153–164. MR 877010 6
- [KS] Tatsuki Kuwagaki and Vivek Shende, *Adjoints, wrapping, and morphisms at infinity*, arXiv:2309.17062. 2
- [KS92] Maximilian Kreuzer and Harald Skarke, *On the classification of quasihomogeneous functions*, *Comm. Math. Phys.* **150** (1992), no. 1, 137–147. MR 1188500 (93k:32075) 5
- [LU21] Yankı Lekili and Kazushi Ueda, *Homological mirror symmetry for Milnor fibers of simple singularities*, *Algebr. Geom.* **8** (2021), no. 5, 562–586. MR 4371540 1, 14
- [LU22] Yankı Lekili and Kazushi Ueda, *Homological mirror symmetry for Milnor fibers via moduli of  $A_\infty$ -structures*, *J. Topol.* **15** (2022), no. 3, 1058–1106. MR 4442683 1, 2, 3, 5, 11, 13
- [McL16] Mark McLean, *Reeb orbits and the minimal discrepancy of an isolated singularity*, *Invent. Math.* **204** (2016), no. 2, 505–594. MR 3489704 14
- [Orl04] D. O. Orlov, *Triangulated categories of singularities and D-branes in Landau-Ginzburg models*, *Tr. Mat. Inst. Steklova* **246** (2004), no. *Algebr. Geom. Metody, Svyazi i Prilozh.*, 240–262. MR MR2101296 1
- [Orl11] Dmitri Orlov, *Formal completions and idempotent completions of triangulated categories of singularities*, *Adv. Math.* **226** (2011), no. 1, 206–217. MR 2735755 6
- [Pet] Chris Peters, *On isolated hypersurface singularities: algebra-geometric and symplectic aspects*, arXiv:2405.03475. 16
- [Pos] Alexander Postnikov, *Total positivity, Grassmannians, and networks*, arXiv:math/0609764. 1
- [Rei83] Miles Reid, *Minimal models of canonical 3-folds*, *Algebraic varieties and analytic varieties* (Tokyo, 1981), *Adv. Stud. Pure Math.*, vol. 1, North-Holland, Amsterdam, 1983, pp. 131–180. MR 715649 14
- [Sco06] Joshua S. Scott, *Grassmannians and cluster algebras*, *Proc. London Math. Soc.* (3) **92** (2006), no. 2, 345–380. MR 2205721 (2007e:14078) 1
- [Seg13] Ed Segal, *The closed state space of affine Landau-Ginzburg B-models*, *J. Noncommut. Geom.* **7** (2013), no. 3, 857–883. MR 3108698 13
- [Sei00] Paul Seidel, *Graded Lagrangian submanifolds*, *Bull. Soc. Math. France* **128** (2000), no. 1, 103–149. MR MR1765826 (2001c:53114) 12
- [Sei08] ———,  *$A_\infty$ -subalgebras and natural transformations*, *Homology, Homotopy Appl.* **10** (2008), no. 2, 83–114. MR 2426130 (2010k:53154) 11
- [Sei09] ———, *Symplectic homology as Hochschild homology*, *Algebraic geometry—Seattle 2005. Part 1*, *Proc. Sympos. Pure Math.*, vol. 80, Amer. Math. Soc., Providence, RI, 2009, pp. 415–434. MR 2483942 (2010c:53129) 12
- [Sei10] ———, *Suspending Lefschetz fibrations, with an application to local mirror symmetry*, *Comm. Math. Phys.* **297** (2010), no. 2, 515–528. MR 2651908 4
- [Sei11] ———, *Homological mirror symmetry for the genus two curve*, *J. Algebraic Geom.* **20** (2011), no. 4, 727–769. MR 2819674 (2012f:53186) 7
- [She11] Nick Sheridan, *On the homological mirror symmetry conjecture for pairs of pants*, *J. Differential Geom.* **89** (2011), no. 2, 271–367. MR 2863919 (2012m:53196) 7, 10
- [She15] ———, *Homological mirror symmetry for Calabi-Yau hypersurfaces in projective space*, *Invent. Math.* **199** (2015), no. 1, 1–186. MR 3294958 8, 10
- [She16] ———, *On the Fukaya category of a Fano hypersurface in projective space*, *Publ. Math. Inst. Hautes Études Sci.* **124** (2016), 165–317. MR 3578916 10
- [She20] ———, *Versality of the relative Fukaya category*, *Geom. Topol.* **24** (2020), no. 2, 747–884. MR 4153652 9

- [SS21] Nick Sheridan and Ivan Smith, *Homological mirror symmetry for generalized Greene-Plesser mirrors*, *Invent. Math.* **224** (2021), no. 2, 627–682. MR 4243021 8
- [Tak10] Atsushi Takahashi, *Weighted projective lines associated to regular systems of weights of dual type*, *New developments in algebraic geometry, integrable systems and mirror symmetry* (RIMS, Kyoto, 2008), *Adv. Stud. Pure Math.*, vol. 59, Math. Soc. Japan, Tokyo, 2010, pp. 371–388. MR 2683215 3
- [Ued06] Kazushi Ueda, *Homological mirror symmetry and simple elliptic singularities*, *math.AG/0604361*, 2006. 3

DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE LONDON, SOUTH KENSINGTON, LONDON, SW7 2AZ  
*Email address:* y.lekili@imperial.ac.uk

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, 3-8-1 KOMABA, MEGURO-KU,  
TOKYO, 153-8914, JAPAN  
*Email address:* kazushi@ms.u-tokyo.ac.jp